

At the level of the cell, good selection has a name. Metabolic flexibility — the capacity to switch fuels as demand changes — is what state agility looks like in the tissues, and its loss is where much of chronic disease begins.

What Metabolic Flexibility Is


The abstract idea of energy-state selection has a concrete cellular expression, and it is one of the most important concepts in modern metabolic health: metabolic flexibility. It is the capacity of the body to switch between fuels — to burn fat when fat is abundant and demand is low, to burn carbohydrate when intensity is high, to shift toward ketones during fasting — matching fuel selection to the changing supply of substrate and demand for energy (Goodpaster & Sparks, 2017). A flexible metabolism reads its situation and selects the appropriate fuel; an inflexible one is stuck.

This is, quite literally, energy-state selection at the cellular scale. The whole-body states of this series — mobilization, assimilation, recovery — each call for a particular pattern of fuel use, and metabolic flexibility is the cellular machinery that delivers the right fuel for each. The fed-to-fasted transition, the rest-to-exercise transition, the shift from a carbohydrate-rich to a fat-rich meal: each is a state change that a flexible metabolism executes smoothly and an inflexible one performs poorly. The cell, in miniature, faces the same selection problem the organism faces.

Seeing metabolic flexibility this way unifies it with the master skill. It is not a separate metabolic curiosity but the tissue-level face of the same competence — the capacity to select and switch appropriately — that the series has argued is the target of health. When we speak of a person becoming better at selecting energy states, part of what we mean, at the cellular level, is that their metabolism has become more flexible, more able to match fuel to demand across the situations of a day.

The concept also gives the abstract idea of selection an objective, measurable anchor. Metabolic flexibility can be assessed — through the respiratory exchange ratio during controlled feeding and exercise, through the glucose response to a meal, through the capacity to shift substrate use under challenge. This makes the cellular face of selection something that can be tracked and improved, not merely described, which is essential for a program that insists on measurement. The glucose curve of a continuous monitor, for instance, is in part a readout of this flexibility.

It is worth pausing on how counterintuitive this is against the usual language of metabolism. We tend to describe good metabolic health in terms of levels — a normal fasting glucose, a healthy lipid panel — as though the goal were to hold certain values steady. Metabolic flexibility reframes the goal as the capacity to change: to move fuels appropriately as the situation shifts, spiking and clearing cleanly rather than holding any single value fixed. Health, here too, lives in the dynamics and the switching



rather than in the steadiness, which is precisely the reframing this series has urged at every level, from the whole-body states down to the individual cell.

A Multi-Organ Competence

Metabolic flexibility was once thought to be mainly a property of skeletal muscle, the tissue whose fuel use shifts most dramatically between rest and exercise. That view has broadened decisively. Contemporary work recognizes metabolic flexibility as a systemic, multi-organ competence, expressed across skeletal muscle, liver, heart, brain, and adipose tissue, each of which must switch fuels appropriately and all of which must coordinate their switching for the organism as a whole to remain flexible (Perspectives on metabolic flexibility, 2025).

This multi-organ character matters because it means metabolic flexibility is a whole-body signature, not a local one. The liver's capacity to switch between storing and releasing glucose, the heart's ability to burn whatever fuel is available, the brain's use of ketones during fasting, and adipose tissue's regulated release of fatty acids all participate in a coordinated system. Flexibility is the property of that coordinated whole, and its loss in any major node — insulin resistance in muscle, fat accumulation in the liver — degrades the flexibility of the entire system.

The systemic view aligns metabolic flexibility precisely with the whole-body energy-state selection this series describes. Just as the organism selects among mobilization, assimilation, and recovery by coordinating many systems at once, metabolic flexibility is the coordinated fuel-switching of many organs, and the two are deeply linked: the whole-body state a person selects sets the fuel demand that their organs must flexibly meet. Metabolic flexibility is, in effect, the metabolic execution of the state the brain has selected.

Understanding flexibility as multi-organ also explains why it is such a sensitive barometer of health. A competence distributed across every major metabolic tissue, and requiring their coordination, is vulnerable to disruption at many points, and its degradation reflects trouble somewhere in the system before that trouble has produced a named disease. This is what makes metabolic flexibility, and its measurable signatures, an early and informative window on the state of a person's regulatory economy — a theme the diagnostics article develops.

Inflexibility as Disease

If flexibility is the cellular signature of health, then metabolic inflexibility is the cellular signature of disease, and the connection is now well established. The core defect of insulin resistance can be described as a metabolic inflexibility: a failure to switch appropriately between fuels, so that the tissues do not shift to burning glucose when insulin signals abundance, nor cleanly to fat when it signals scarcity. The mismatch between fuel availability and fuel oxidation is a defining feature of the metabolically unwell state (Metabolic flexibility and inflexibility, 2024).

This inflexibility appears early in the trajectory toward cardiometabolic disease, often before conventional diagnostic thresholds are crossed. A metabolism that has begun to lose its flexibility — spiking and lingering after a meal, failing to draw down fat between meals — is a metabolism heading toward trouble, and it can be detected in the dynamics of its fuel handling well before a fasting glucose or an HbA1c would flag it. Inflexibility is, in a sense, the functional prodrome of metabolic disease, visible in how the system switches long before it is visible in a static value.

Read through this series' framework, metabolic inflexibility is being stuck at the cellular level — the fuel-switching equivalent of the stuck whole-body states of Article 2. A metabolism locked onto one fuel, unable to shift with demand, is in the same predicament as an organism locked in mobilization or unable to enter



recovery: it has lost the capacity for context-appropriate change that defines health. The same pathology of rigidity that afflicts the whole-body selector afflicts the cellular one, and for the same underlying reason.

This parallel is more than an analogy; the two levels are causally linked. A person chronically stuck in the wrong whole-body states — perpetually stressed, poorly recovered, metabolically overloaded — imposes on their cells a pattern of fuel demand that erodes cellular flexibility, while cellular inflexibility in turn constrains the whole-body states the person can effectively enter. Rigidity at one level feeds rigidity at the other, which is part of why the stuck states of chronic disease are so self-reinforcing, and why restoring flexibility often requires acting at both levels at once.

Mitochondria as the Switch

The physical machinery of metabolic flexibility is, in large part, the mitochondrion. It is in the mitochondria that fuels are actually oxidized, and the capacity to switch cleanly between burning fat and carbohydrate depends on the number, quality, and adaptability of a cell's mitochondrial network. A robust network switches fuels readily; a depleted or damaged one becomes rigid, unable to accommodate shifting demand. The mitochondrion is the cellular switch on which flexibility ultimately runs (Picard & Shirihai, 2022).

This connects metabolic flexibility directly to the mitochondrial adaptations developed in the mobilization series. The same exercise-driven mitochondrial biogenesis and quality control that build endurance also build the substrate of metabolic flexibility, which is why physical training so reliably improves the capacity to switch fuels. Mitohormesis — the adaptive response to the transient stress of exertion — expands and cleans the network that flexibility depends on, tying the flexibility of the cell to the deliberate selection of the active state (Gunawan et al., 2025).

The mitochondrial basis of flexibility also explains its links to aging and to the other energy states. Mitochondrial quality is a shared substrate across mobilization, assimilation, and recovery: the network that exercise builds is the one that recovery repairs and that digestion fuels, and its condition determines the flexibility available to all three states. An improvement in mitochondrial health through any one pillar enhances metabolic flexibility across the board, which is one mechanism by which the pillars compound.

Because the mitochondrion is the switch, its decline is felt as a loss of flexibility. Damaged, inefficient mitochondria leak more reactive species, waste more energy, and switch fuels less cleanly, and their accumulation — a recognized hallmark of aging — manifests as the metabolic rigidity of older tissue (López-Otín et al., 2023). Conversely, interventions that preserve or restore mitochondrial quality restore flexibility, which is why the mitochondrion sits at the center of both the mechanism and the treatment of metabolic inflexibility.

Flexibility Declines with Age — and Responds to Training

Metabolic flexibility, like the other capacities in this series, tends to decline across the lifespan. Older metabolisms switch fuels less readily, clear meals more sluggishly, and draw down fat less efficiently, part of the broad loss of adaptive capacity that characterizes aging. This decline is not merely cosmetic; it underlies much of the rising cardiometabolic risk of later life, and it is a concrete instance of the loss of selection capacity — cellular stuckness accumulating with the years.

Crucially, though, metabolic flexibility is trainable, and it responds to the same behavioral levers that build the other capacities. Exercise, by building the mitochondrial network, is among the most powerful stimuli for restoring flexibility. Periods of fasting and the deliberate variation of fuel availability challenge the switching machinery and improve it. Attention to the timing of meals aligns fuel demand with the body's rhythms and



enhances flexible handling (Chrononutrition and circadian rhythms, 2025). The capacity to switch fuels, like the capacity to select states, can be built.

The trainability of metabolic flexibility is visible in its dynamic signatures. As a person becomes more flexible through training and appropriate feeding, their glucose excursions after meals shrink and clear faster, their capacity to burn fat at rest improves, and their metabolism handles the transitions of the day more smoothly — changes readable in the continuous glucose traces that have made metabolic dynamics visible outside the laboratory. The improvement can be measured, which means it can be pursued and confirmed rather than merely assumed.

This trainability is what makes metabolic flexibility a target rather than merely a marker. It is not a fixed feature of a person's constitution but a capacity they can build and preserve through the deliberate selection of the active state, the appropriate timing of feeding, and the recovery that maintains the mitochondrial substrate. In the language of the program, improving metabolic flexibility is improving the cellular face of energy-state selection — one of the concrete things that optimizing the master skill actually accomplishes.

The Metabolic Half of the Picture

Metabolic flexibility is the cellular answer to one part of the selection problem: which fuel. It determines whether the tissues can supply the right substrate for whatever state the body enters, switching cleanly between fat and carbohydrate as mobilization, assimilation, and recovery each demand. It is the metabolic execution of state selection, and its presence or absence is much of what separates a body that can enter its states effectively from one that cannot.

But which fuel is only half of the picture. The other half is which branch and how fast — the rapid, whole-body autonomic switching that shifts the organism between sympathetic and parasympathetic configurations, sets the pace of transitions, and determines whether the body can change states in seconds rather than minutes. Metabolic flexibility supplies the fuel appropriate to a state; autonomic flexibility executes the shift into and out of the state itself. The two are complementary faces of the same underlying competence.

Together they constitute what flexible selection looks like in the tissues: a metabolism that switches fuels appropriately and a nervous system that switches states rapidly and reversibly. A person strong in both can select and execute their energy states with precision; a person weak in either is constrained, however good their intentions. The two flexibilities are the physiological substance of the master skill, and both are measurable, trainable, and predictive of health.

The next article turns to the autonomic half — the fast selector that reconfigures the whole body between states, indexed by heart rate variability, and central to both the aging of the organism and the levers by which selection can be improved. With the cellular and systemic faces of flexibility both established, the series will have the physiological foundation it needs to turn to the individuality of selection and, ultimately, to its measurement and optimization.

References

- **Goodpaster BH, Sparks LM.** (2017). Metabolic flexibility in health and disease. *Cell Metabolism*, 25(5), 1027–1036.
- Perspectives on whole-body and tissue-specific metabolic flexibility and implications in cardiometabolic diseases. (2025). *Cell Reports Medicine*.
- Metabolic flexibility and inflexibility: pathology underlying metabolism dysfunction. (2024). Review.
- **Picard M, Shirihaï OS.** (2022). Mitochondrial signal transduction. *Cell Metabolism*, 34(11), 1620–1653.



- **Gunawan AL, Liparulo I, Stahl A.** (2025). Mammalian mitohormesis: from mitochondrial stressors to organismal benefits. *The EMBO Journal*, 44(20), 5640–5661.
- **Memme JM, Erlich AT, Phukan G, Hood DA.** (2021). Exercise and mitochondrial health. *Journal of Physiology*, 599(3), 803–817.
- **López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G.** (2023). Hallmarks of aging: an expanding universe. *Cell*, 186(2), 243–278.
- Chrononutrition and energy balance: how meal timing and circadian rhythms shape weight regulation and metabolic health. (2025). *Nutrients*, 17(13), 2135.
- Glycaemic variability, assessed with continuous glucose monitors, is associated with diet, lifestyle and health in people without diabetes. (2023). *Diabetologia* / related.
- **Bobba-Alves N, Juster R-P, Picard M.** (2022). The energetic cost of allostasis and allostatic load. *Psychoneuroendocrinology*, 146, 105951.
- **Sterling P.** (2012). Allostasis: a model of predictive regulation. *Physiology & Behavior*, 106(1), 5–15.
- **Muoio DM.** (2014). Metabolic inflexibility: when mitochondrial indecision leads to metabolic gridlock. *Cell*, 159(6), 1253–1262.
- Foundational BioSystems. Exercise Is Mitochondrial Training; Core Themes. Internal white papers.

